

Ddx54 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11203**Specification**

Ddx54 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8K4L0
Other Accession	NM_028041 , NP_082317
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Mouse, Rabbit, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98kDa KDa

Ddx54 antibody - C-terminal region - Additional Information**Gene ID** 71990**Alias Symbol** 2410015A15Rik, AI414901, DP97**Other Names**

ATP-dependent RNA helicase DDX54, 3.6.4.13, DEAD box protein 54, Ddx54

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ddx54 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ddx54 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ddx54 antibody - C-terminal region - Protein Information**Name** Ddx54**Function**

Has RNA-dependent ATPase activity. Represses the transcriptional activity of nuclear receptors (By similarity).

Cellular Location


Nucleus, nucleolus.

Ddx54 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ddx54 antibody - C-terminal region - Images



WB Suggested Anti-Ddx54 Antibody Titration: 0.2-1 µg/ml
Positive Control: SP2/0 cell lysate