

Jundm2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11211**Specification**

Jundm2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P97875
Other Accession	NM_030887 , NP_112149
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19kDa KDa

Jundm2 antibody - C-terminal region - Additional Information**Gene ID** 81703

Alias Symbol	Jdp2, Jundp2, TIF, Jundm2
Other Names	
Jun dimerization protein 2, Jdp2, Jundm2	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Jundm2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Jundm2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Jundm2 antibody - C-terminal region - Protein Information**Name** Jdp2**Synonyms** Jundm2**Function**

Component of the AP-1 transcription factor that represses transactivation mediated by the Jun family of proteins. Involved in a variety of transcriptional responses associated with AP-1, such as UV- induced apoptosis, cell differentiation, tumorigenesis and antitumogeneris. Can also function as a repressor by recruiting histone deacetylase 3/HDAC3 to the promoter region of JUN. May control transcription via direct regulation of the modification of histones and the assembly of chromatin (By similarity).

Cellular Location

Nucleus.


Tissue Location

Ubiquitously expressed in all adult tissues tested as well in embryos.

Jundm2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Jundm2 antibody - C-terminal region - Images

WB Suggested Anti-Jundm2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Heart