

Dnajc17 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11229**Specification**

Dnajc17 antibody - middle region - Product Information

Application	WB
Primary Accession	O91WT4
Other Accession	NM_139139 , NP_631878
Reactivity	Human, Mouse, Rat, Horse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

Dnajc17 antibody - middle region - Additional Information**Gene ID** 69408**Alias Symbol** 1700025B16Rik, C87112, D9Bwg1371e**Other Names**

Dnaj homolog subfamily C member 17, Dnajc17

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Dnajc17 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Dnajc17 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Dnajc17 antibody - middle region - Protein Information**Name** Dnajc17**Function**

May negatively affect PAX8-induced thyroglobulin/TG transcription.

Cellular LocationCytoplasm {ECO:0000250|UniProtKB:D3ZSC8}. Nucleus {ECO:0000250|UniProtKB:D3ZSC8}.
Note=Predominantly nuclear {ECO:0000250|UniProtKB:D3ZSC8}**Tissue Location**


Expressed in the thyroid gland.

Dnajc17 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dnajc17 antibody - middle region - Images



WB Suggested Anti-Dnajc17 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Muscle