

Duxbl antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11274**Specification**

Duxbl antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q7TNE6
Other Accession	NM_183389 , NP_899245
Reactivity	Mouse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39kDa KDa

Duxbl antibody - C-terminal region - Additional Information

Alias Symbol **Duxl, 1110051B16Rik, Duxbl**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Duxbl antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Duxbl antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Duxbl antibody - C-terminal region - Protein Information**Duxbl antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Duxbl antibody - C-terminal region - Images



WB Suggested Anti-Duxbl Antibody Titration: 0.125µg/ml

ELISA Titer: 1:1562500

Positive Control: SP2/0 cell lysate

Duxbl antibody - C-terminal region - References

Katayama, S., et al., (2005) Science 309 (5740), 1564-1566
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.