

Mafa antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11277**Specification**

Mafa antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8°CF90
Other Accession	NM_194350 , NP_919331
Reactivity	Human, Mouse, Rat
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37kDa KDa

Mafa antibody - C-terminal region - Additional Information

Alias Symbol **RIPE3b1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Mafa antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


Mafa antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Mafa antibody - C-terminal region - Protein Information**Mafa antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mafa antibody - C-terminal region - Images



90 kDa
65 kDa
40 kDa
31 kDa
22 kDa

WB Suggested Anti-Mafa Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Mouse Muscle