

Naca-rs Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11340**Specification**

Naca-rs Antibody - C-terminal region - Product Information

Application	WB
Other Accession	XM_354689 , XP_354689
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

Naca-rs Antibody - C-terminal region - Additional Information

Alias Symbol **Gm1878**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Naca-rs antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


Naca-rs Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Naca-rs Antibody - C-terminal region - Protein Information**Naca-rs Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Naca-rs Antibody - C-terminal region - Images



Host: Rabbit
Target Name: Naca-rs
Sample Tissue: Mouse Stomach lysates
Antibody Dilution: 1.0µg/ml