

Vgll1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11341**Specification**

Vgll1 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O99NC0
Other Accession	NM_133251 , NP_573514
Reactivity	Mouse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

Vgll1 Antibody - C-terminal region - Additional Information**Gene ID** 170828**Alias Symbol** **Tondu****Other Names**

Transcription cofactor vestigial-like protein 1, Vgl-1, Vestigial-related factor, Vgll1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Vgll1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Vgll1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Vgll1 Antibody - C-terminal region - Protein Information**Name** Vgll1**Function**

May act as a specific coactivator for the mammalian TEFs.

Cellular Location


Nucleus.

Vgll1 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Vgll1 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: Vgll1
Sample Tissue: Mouse Stomach lysates
Antibody Dilution: 1.0µg/ml