

Hpcal1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11362**Specification**

Hpcal1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P62748
Other Accession	NM_016677 , NP_057886
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Sheep, Horse, Bovine, Dog
Predicted Host	Human, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 22kDa KDa

Hpcal1 antibody - C-terminal region - Additional Information**Gene ID** 53602**Alias Symbol** NVP-3, Nvp3, VILIP3, VnsI3, VsnI3**Other Names**

Hippocalcin-like protein 1, Neural visinin-like protein 3, NVL-3, NVP-3, Visinin-like protein 3, VILIP-3, Hpcal1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hpcal1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Hpcal1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Hpcal1 antibody - C-terminal region - Protein Information**Name** Hpcal1**Function**

May be involved in the calcium-dependent regulation of rhodopsin phosphorylation.

Cellular Location


Membrane {ECO:0000250|UniProtKB:P37235}; Lipid- anchor {ECO:0000250|UniProtKB:P37235}

Hpcal1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hpcal1 antibody - C-terminal region - Images



WB Suggested Anti-Hpcal1 Antibody Titration: 1.0 μ g/ml
Positive Control: Mouse Heart