

HTLF antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11363**Specification**

HTLF antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q6P4Q2
Other Accession	NM_002158 , NP_002149
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Chicken, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 47kDa KDa

HTLF antibody - N-terminal region - Additional Information

Alias Symbol **HTLF**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HTLF antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


HTLF antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HTLF antibody - N-terminal region - Protein Information**HTLF antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HTLF antibody - N-terminal region - Images



WB Suggested Anti-HTLF Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

HTLF antibody - N-terminal region - References

Li,C., (1992) Genomics 13 (3), 658-664
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.