

SMARCD2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11379**Specification**

SMARCD2 antibody - C-terminal region - Product Information

Application	WB, IHC
Primary Accession	O92925-2
Other Accession	NM_003077 , NP_003068
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rabbit, Zebrafish, Pig, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

SMARCD2 antibody - C-terminal region - Additional Information

Alias Symbol **BAF60B, CRACD2, PRO2451, Rsc6p**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SMARCD2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


SMARCD2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SMARCD2 antibody - C-terminal region - Protein Information**SMARCD2 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SMARCD2 antibody - C-terminal region - Images



Rabbit Anti-SMARCD2 Antibody
Paraffin Embedded Tissue: Human Lung
Cellular Data: Alveolar cells
Antibody Concentration: 16 µg/ml
Magnification: 400X



Rabbit Anti-SMARCD2 Antibody
Paraffin Embedded Tissue: Human cardiac cell
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-SMARCD2 Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate
SMARCD2 is supported by BioGPS gene expression data to be expressed in Jurkat

SMARCD2 antibody - C-terminal region - References

Lehner,B. (2004) Genome Res. 14 (7), 1315-1323 Reconstitution and Storage:For short term use,

store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.