

TEAD4 Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11384**Specification**

TEAD4 Antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	O15561-3
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44kDa KDa

TEAD4 Antibody - N-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TEAD4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


TEAD4 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TEAD4 Antibody - N-terminal region - Protein Information**TEAD4 Antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TEAD4 Antibody - N-terminal region - Images



Rabbit Anti-TEAD4 Antibody

Catalog Number: AI11384

Formalin Fixed Paraffin Embedded Tissue: Human Adult liver Observed Staining: Nuclear in endothelial cells not in hepatocytes

Primary Antibody

Concentration: 1:600

Secondary Antibody: Donkey anti-Rabbit-Cy2/3


Secondary Antibody

Concentration: 1:200

Magnification: 20X

Exposure Time: 0.5 – 2.0 sec

Protocol located in Reviews and Data.



90 kDa
65 kDa
40 kDa
29 kDa
22 kDa

Host: Rabbit

Target Name: TEAD4

Sample Tissue: Thyroid Tumor lysates

Antibody Dilution: 1.0 µg/ml



250 kDa
150 kDa
100 kDa
75 kDa
50 kDa
37 kDa
25 kDa
20 kDa
15 kDa
10 kDa

WB Suggested Anti-TEAD4 antibody Titration: 1 µg/ml

Sample Type: Human heart