

NR2C1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11386**Specification**

NR2C1 antibody - N-terminal region - Product Information

Application	WB, IHC
Primary Accession	Q15626
Other Accession	NM_001032287 , NP_001027458
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

NR2C1 antibody - N-terminal region - Additional Information

Alias Symbol **TR2, TR2-11**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NR2C1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NR2C1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NR2C1 antibody - N-terminal region - Protein Information**NR2C1 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR2C1 antibody - N-terminal region - Images



Immunohistochemistry with Human Skin lysate tissue at an antibody concentration of 5.0 µg/ml using anti-NR2C1 antibody (AI11386)



WB Suggested Anti-NR2C1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: 721_B cell lysate

NR2C1 antibody - N-terminal region - References

Lin, Y.L., (2006) Biochem. Biophys. Res. Commun. 350 (2), 430-436 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.