

NAB2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11449**Specification**

NAB2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q15742
Other Accession	NM_005967 , NP_005958
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Rabbit, Pig, Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 57kDa KDa

NAB2 antibody - N-terminal region - Additional Information**Gene ID** 4665

Alias Symbol	MADER
--------------	--------------

Other Names

NGFI-A-binding protein 2, EGR-1-binding protein 2, Melanoma-associated delayed early response protein, Protein MADER, NAB2, MADER

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-NAB2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

NAB2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NAB2 antibody - N-terminal region - Protein Information**Name** NAB2**Synonyms** MADER**Function**

Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability (By similarity).

Cellular Location

Nucleus. Note=Isoform 2 is not localized to the nucleus.


Tissue Location

Widely expressed at low levels. Highly expressed in melanoma cell lines

NAB2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NAB2 antibody - N-terminal region - Images

WB Suggested Anti-NAB2 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:1562500

Positive Control: HepG2 cell lysate

NAB2 antibody - N-terminal region - References

Srinivasan,R., et al., (er) J. Biol. Chem. (2006) In press
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.