

NR1H2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11474**Specification**

NR1H2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q68°CY8
Other Accession	NM_007121 , NP_009052
Reactivity	Human, Rat, Horse, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

NR1H2 antibody - N-terminal region - Additional Information

Alias Symbol **LXR-b, LXRB, NER, NER-I, RIP15, UNR**
Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-NR1H2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


NR1H2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NR1H2 antibody - N-terminal region - Protein Information**NR1H2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR1H2 antibody - N-terminal region - Images



70 kDa
60 kDa
48 kDa
36 kDa
21 kDa

WB Suggested Anti-NR1H2 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:62500

Positive Control: Transfected 293T

NR1H2 antibody - N-terminal region - References

Albers, M., (2006) J. Biol. Chem. 281 (8), 4920-4930 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. Publications: Bogan, R. L. & Hennebold, J. D. The reverse cholesterol transport system as a potential mediator of luteolysis in the primate corpus luteum. Reproduction 139, 163-76 (2010). WB, IHC, Human, Dog, Rat, Guinea pig, H19776099