

Ppargc1a antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11479

Specification

Ppargc1a antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O70343
Other Accession	NM_008904 , NP_032930
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Goat, Horse, Bovine, Dog
Predicted	Human, Mouse, Rabbit, Zebrafish, Pig, Chicken, Goat, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	90kDa KDa

Ppargc1a antibody - C-terminal region - Additional Information

Gene ID 19017

Alias Symbol A830037N07Rik, PGC-1, PGC-1v, Pgc-1alpha, Pgc1, Pgco1, Ppargc1, Gm11133, ENSMUSG00000079510

Other Names

Peroxisome proliferator-activated receptor gamma coactivator 1-alpha, PGC-1-alpha, PPAR-gamma coactivator 1-alpha, PPARGC-1-alpha, Ppargc1a, Pgc1, Pgc1a, Ppargc1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ppargc1a antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ppargc1a antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ppargc1a antibody - C-terminal region - Protein Information

Name Ppargc1a

Synonyms Pgc1, Pgc1a, Ppargc1

Function

Transcriptional coactivator for steroid receptors and nuclear receptors (PubMed:15744310, PubMed:<a

[12754525](http://www.uniprot.org/citations/12754525), PubMed: [23217713](http://www.uniprot.org/citations/23217713), PubMed: [9529258](http://www.uniprot.org/citations/9529258)). Greatly increases the transcriptional activity of PPARG and thyroid hormone receptor on the uncoupling protein promoter (PubMed: [15744310](http://www.uniprot.org/citations/15744310), PubMed: [12754525](http://www.uniprot.org/citations/12754525), PubMed: [23217713](http://www.uniprot.org/citations/23217713), PubMed: [9529258](http://www.uniprot.org/citations/9529258)). Can regulate key mitochondrial genes that contribute to the program of adaptive thermogenesis (PubMed: [15744310](http://www.uniprot.org/citations/15744310), PubMed: [12754525](http://www.uniprot.org/citations/12754525), PubMed: [23217713](http://www.uniprot.org/citations/23217713), PubMed: [9529258](http://www.uniprot.org/citations/9529258)). Plays an essential role in metabolic reprogramming in response to dietary availability through coordination of the expression of a wide array of genes involved in glucose and fatty acid metabolism (PubMed: [15744310](http://www.uniprot.org/citations/15744310), PubMed: [12754525](http://www.uniprot.org/citations/12754525), PubMed: [23217713](http://www.uniprot.org/citations/23217713), PubMed: [9529258](http://www.uniprot.org/citations/9529258)). Acts as a key regulator of gluconeogenesis: stimulates hepatic gluconeogenesis by increasing the expression of gluconeogenic enzymes, and acting together with FOXO1 to promote the fasting gluconeogenic program (PubMed: [12754525](http://www.uniprot.org/citations/12754525)). Induces the expression of PERM1 in the skeletal muscle in an ESRRRA-dependent manner (By similarity). Also involved in the integration of the circadian rhythms and energy metabolism (PubMed: [17476214](http://www.uniprot.org/citations/17476214)). Required for oscillatory expression of clock genes, such as BMAL1 and NR1D1, through the coactivation of RORA and RORC, and metabolic genes, such as PDK4 and PEPCK (PubMed: [17476214](http://www.uniprot.org/citations/17476214)).

Cellular Location

Nucleus. Nucleus, PML body

Tissue Location


White quadriceps and red tibialis anterior (TA) muscles, liver, kidney and brown adipose tissue (at protein level) Skeletal muscle, brown adipose tissue, heart, kidney and brain


Ppargc1a antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ppargc1a antibody - C-terminal region - Images





168 kDa
144 kDa
90 kDa
65 kDa
40 kDa

WB Suggested Anti-Ppargc1a Antibody Titration: 0.2-1 $\mu\text{g/ml}$

ELISA Titer: 1:312500

Positive Control: Mouse Brain