

UHRF1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11480**Specification**

UHRF1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q2HIX7
Other Accession	NM_013282 , NP_037414
Reactivity	Human, Mouse, Rat, Zebrafish, Bovine, Dog
Predicted	Mouse, Rat, Zebrafish, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	91kDa KDa

UHRF1 antibody - N-terminal region - Additional Information

Alias Symbol	FLJ21925, ICBP90, MGC138707, Np95, RNF106, hNP95
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-UHRF1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


UHRF1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

UHRF1 antibody - N-terminal region - Protein Information**UHRF1 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UHRF1 antibody - N-terminal region - Images



WB Suggested Anti-UHRF1 Antibody Titration: 0.2-1 μ g/ml
ELISA Titer: 1:312500
Positive Control: Human Stomach



WB Suggested Anti-UHRF1 Antibody
Positive Control: Lane 1: 60ug HCT116 lysate Lane 2: 60ug HCT116 lysate + siRNA
Primary Antibody Dilution : 1:1000
Secondary Antibody : Anti rabbit-HRP
Secondary Antibody Dilution : 1:5,000
Submitted by: Chinweike Ukomadu, Brigham and Women's Hospital, Boston

UHRF1 antibody - N-terminal region - References

Achour,M., (2008) Oncogene 27 (15), 2187-2197 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.