

**ZNF777 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI11507****Specification**

---

**ZNF777 antibody - C-terminal region - Product Information**

|                   |                                                      |
|-------------------|------------------------------------------------------|
| Application       | WB                                                   |
| Primary Accession | <a href="#">Q8N2J5</a>                               |
| Other Accession   | <a href="#">NM_015694</a> , <a href="#">EAL24432</a> |
| Reactivity        | Human                                                |
| Predicted         | Human                                                |
| Host              | Rabbit                                               |
| Clonality         | Polyclonal                                           |
| Calculated MW     | 58kDa KDa                                            |

**ZNF777 antibody - C-terminal region - Additional Information****Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-ZNF777 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


**Precautions**

ZNF777 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**ZNF777 antibody - C-terminal region - Protein Information****ZNF777 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ZNF777 antibody - C-terminal region - Images**



WB Suggested Anti-ZNF777 Antibody Titration: 0.2-1 µg/ml

Positive Control: Jurkat cell lysate

### **ZNF777 antibody - C-terminal region - References**

Nagase T, (1999) DNA Res. 1999 Oct 29;6(5):337-45. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.