

**ZNF70 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI11568****Specification**

---

**ZNF70 antibody - C-terminal region - Product Information**

|                   |                                                               |
|-------------------|---------------------------------------------------------------|
| Application       | WB, IHC                                                       |
| Primary Accession | <a href="#">O9UC06</a>                                        |
| Other Accession   | <a href="#">NM_021916</a> , <a href="#">NP_068735</a>         |
| Reactivity        | Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog |
| Predicted Host    | Human, Horse, Bovine                                          |
| Clonality         | Rabbit                                                        |
| Calculated MW     | Polyclonal<br>51kDa KDa                                       |

**ZNF70 antibody - C-terminal region - Additional Information****Gene ID** 7621

|                                                            |              |
|------------------------------------------------------------|--------------|
| Alias Symbol                                               | <b>Cos17</b> |
| <b>Other Names</b>                                         |              |
| Zinc finger protein 70, Zinc finger protein N27C7-1, ZNF70 |              |

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-ZNF70 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

ZNF70 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**ZNF70 antibody - C-terminal region - Protein Information****Name** ZNF70**Function**

May be involved in transcriptional regulation.

**Cellular Location**


Nucleus.

**ZNF70 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ZNF70 antibody - C-terminal region - Images**



Human Stomach



WB Suggested Anti-ZNF70 Antibody Titration: 5.0µg/ml

ELISA Titer: 1:312500

Positive Control: HepG2 cell lysate

#### **ZNF70 antibody - C-terminal region - References**

Collins, J.E., Genome Biol. 5 (10), R84 (2004) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.