

ZNF397 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11601**Specification**

ZNF397 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q8NF99-2
Other Accession	NM_032347 , NP_115723
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31kDa KDa

ZNF397 antibody - C-terminal region - Additional Information

Alias Symbol **ZNF47, ZSCAN15**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZNF397 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


ZNF397 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF397 antibody - C-terminal region - Protein Information**ZNF397 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF397 antibody - C-terminal region - Images



Human Skin



WB Suggested Anti-ZNF397 Antibody Titration: 1.25 μ g/ml
Positive Control: HepG2 cell lysate

ZNF397 antibody - C-terminal region - References

Wu,Y., Gene 310, 193-201 (2003)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.