

INSM2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11604**Specification**

INSM2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O96Q84
Other Accession	NM_032594 , NP_115983
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa KDa

INSM2 antibody - N-terminal region - Additional Information

Alias Symbol **IA6, IA-6, mlt1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-INSM2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


INSM2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

INSM2 antibody - N-terminal region - Protein Information**INSM2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

INSM2 antibody - N-terminal region - Images



WB Suggested Anti-INSM2 Antibody Titration: 2.5µg/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

INSM2 antibody - N-terminal region - References

Ota, T., (2004) Nat. Genet. 36 (1), 40-45
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.