

BTBD6 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11614**Specification**

BTBD6 antibody - middle region - Product Information

Application	WB
Primary Accession	O96KE9
Other Accession	NM_033271 , NP_150374
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Rabbit, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 46kDa KDa

BTBD6 antibody - middle region - Additional Information**Gene ID** 90135

Alias Symbol	BDPL
Other Names	
BTB/POZ domain-containing protein 6, Lens BTB domain protein, BTBD6, BDPL	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-BTBD6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

BTBD6 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

BTBD6 antibody - middle region - Protein Information**Name** BTBD6 ([HGNC:19897](#))**Synonyms** BDPL**Function**

Adapter protein for the cul3 E3 ubiquitin-protein ligase complex (By similarity). Involved in late neuronal development and muscle formation (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q2LE78}. Note=Found in punctated bodies in the cytoplasm {ECO:0000250|UniProtKB:Q2LE78}


Tissue Location

Expressed in lens..

BTBD6 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BTBD6 antibody - middle region - Images

WB Suggested Anti-BTBD6 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:12500

Positive Control: Human brain

BTBD6 antibody - middle region - References

Wistow, G., (et al) Mol. Vis. 8, 171-184 (2002) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.