

ZNF564 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11630**Specification**

ZNF564 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8TBZ8
Other Accession	NM_144976 , NP_659413
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Mouse, Rat, Zebrafish, Pig, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 64kDa KDa

ZNF564 antibody - C-terminal region - Additional Information**Gene ID** 163050

Alias Symbol	MGC26914
Other Names	
Zinc finger protein 564, ZNF564	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF564 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF564 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF564 antibody - C-terminal region - Protein Information**Name** ZNF564**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF564 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF564 antibody - C-terminal region - Images



WB Suggested Anti-ZNF564 Antibody Titration: 0.2-1 µg/ml

Positive Control: Jurkat cell lysate

ZNF564 antibody - C-terminal region - References

Isogai, T. Unpublished (2002) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.