

ZNF498 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11631**Specification**

ZNF498 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q14C99
Other Accession	NM_145115 , AAI14574
Reactivity	Human, Mouse, Rat, Rabbit, Bovine, Dog
Predicted	Human, Rabbit, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

ZNF498 antibody - C-terminal region - Additional Information

Alias Symbol **ZSCAN25, ZNF498**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZNF498 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


ZNF498 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF498 antibody - C-terminal region - Protein Information**ZNF498 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF498 antibody - C-terminal region - Images



WB Suggested Anti-ZNF498 Antibody Titration: 5.0µg/ml

ELISA Titer: 1:312500

Positive Control: Jurkat cell lysate

ZNF498 antibody - C-terminal region - References

Filion GJ,.(2006) Mol Cell Biol. Jan;26(1):169-81.Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.