

DHR SX antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11632**Specification**

DHR SX antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8N5I4
Other Accession	NM_145177 , NP_660160
Reactivity	Human, Rat, Yeast, Bovine
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36kDa KDa

DHR SX antibody - N-terminal region - Additional Information**Gene ID** 207063**Alias Symbol** CXorf11, DHR S5X, DHR S5Y, DHR SXY, DHR SY, SDR46C1**Other Names**

Dehydrogenase/reductase SDR family member on chromosome X, 1.1.-., DHR SXY, DHR SX, CXorf11, DHR S5X

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DHR SX antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

DHR SX antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DHR SX antibody - N-terminal region - Protein Information**Name** DHR SX**Synonyms** CXorf11, DHR S5X, SDR46C1, SDR7C6**Function**

Involved in the positive regulation of starvation-induced autophagy (PubMed:25076851).

Cellular Location

Secreted. Note=Secreted in a non-classical form. A signal peptide sequence at position 1-31 is

predicted


Tissue Location

Widely expressed. Highly expressed in the pancreas.

DHRX antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DHRX antibody - N-terminal region - Images

WB Suggested Anti-DHRX Antibody Titration: 1.0 µg/ml
Positive Control: THP-1 Whole Cell