

ZNF398 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11655**Specification**

ZNF398 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8TD17
Other Accession	NM_170686 , NP_733787
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Rabbit, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 71kDa KDa

ZNF398 antibody - C-terminal region - Additional Information**Gene ID** 57541**Alias Symbol** P51, P71, ZER6**Other Names**

Zinc finger protein 398, Zinc finger DNA-binding protein p52/p71, ZNF398, KIAA1339, ZER6

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF398 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF398 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF398 antibody - C-terminal region - Protein Information**Name** ZNF398**Synonyms** KIAA1339, ZER6**Function**

Functions as a transcriptional activator.

Cellular Location


Nucleus.

ZNF398 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF398 antibody - C-terminal region - Images



WB Suggested Anti-ZNF398 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: HepG2 cell lysate

ZNF398 antibody - C-terminal region - References

Scherer, S.W., (2003) Science 300 (5620), 767-772
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.