

NFATC1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11657**Specification**

NFATC1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q2M1S3
Other Accession	NM_172390 , NP_765978
Reactivity	Human, Mouse, Rat, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78kDa KDa

NFATC1 antibody - N-terminal region - Additional Information

Alias Symbol **MGC138448, NF-ATC, NFAT2, NFATc**
Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NFATC1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


NFATC1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NFATC1 antibody - N-terminal region - Protein Information**NFATC1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NFATC1 antibody - N-terminal region - Images

**Rabbit Anti-NFATC1 Antibody**

Catalog Number: AI11657

Formalin Fixed Paraffin Embedded Tissue: Human Adult heart Observed Staining: Nuclear, Cytoplasmic

Primary Antibody

Concentration: 1:600

Secondary Antibody: Donkey anti-Rabbit-Cy2/3


Secondary Antibody

Concentration: 1:200

Magnification: 20X

Exposure Time: 0.5 – 2.0 sec

Protocol located in Reviews and Data.



WB Suggested Anti-NFATC1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

NFATC1 is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells

NFATC1 antibody - N-terminal region - References

Crotti, T.N., (2008) J. Cell. Physiol. 215 (3), 636-644 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.