

ZNF527 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11670**Specification**

ZNF527 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NB42
Other Accession	XM_030378 , XP_030378
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Human, Rabbit, Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 71kDa KDa

ZNF527 antibody - C-terminal region - Additional Information**Gene ID** 84503**Other Names**

Zinc finger protein 527, ZNF527, KIAA1829

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZNF527 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF527 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF527 antibody - C-terminal region - Protein Information**Name** ZNF527**Synonyms** KIAA1829**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF527 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF527 antibody - C-terminal region - Images



WB Suggested Anti-ZNF527 Antibody Titration: 2.5µg/ml

ELISA Titer: 1:312500

Positive Control: HepG2 cell lysate

ZNF527 antibody - C-terminal region - References

Ota T, (2004) Nat Genet. 36(1):40-5
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.