

TBX20 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11676**Specification**

TBX20 antibody - middle region - Product Information

Application	WB
Primary Accession	Q000T4
Other Accession	NM_001077653 , NP_001071121
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50kDa KDa

TBX20 antibody - middle region - Additional Information

Alias Symbol **ASD4**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TBX20 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


TBX20 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

TBX20 antibody - middle region - Protein Information**TBX20 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TBX20 antibody - middle region - Images



WB Suggested Anti-TBX20 Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate

TBX20 antibody - middle region - References

Kirk, E.P., (2007) Am. J. Hum. Genet. 81 (2), 280-291 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.