

Lcorl antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11679**Specification**

Lcorl antibody - middle region - Product Information

Application	WB
Primary Accession	Q3U285
Other Accession	NM_172153 , NP_742165
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa kDa

Lcorl antibody - middle region - Additional Information**Gene ID** 209707**Alias Symbol** **Mlr1****Other Names**

Ligand-dependent nuclear receptor corepressor-like protein, LCoR-like protein, Mblk1-related protein 1, Transcription factor MLR1, Lcorl, Mlr1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Lcorl antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Lcorl antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Lcorl antibody - middle region - Protein Information**Name** Lcorl**Synonyms** Mlr1**Function**

May act as transcription activator that binds DNA elements with the sequence 5'-CCCTATCGATCGATCTCTACCT-3'. May play a role in spermatogenesis.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00320}.


Tissue Location

Expressed predominantly in the spermatocytes of the testis. Also found in the kidney, liver and heart

Lcorl antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lcorl antibody - middle region - Images

WB Suggested Anti-Lcorl Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Kidney