

BRF1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11681**Specification**

BRF1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8°CFK2
Other Accession	NM_028193 , NP_082469
Reactivity	Mouse, Rat, Rabbit
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74kDa KDa

BRF1 antibody - C-terminal region - Additional Information

Alias Symbol	GTF3B, TAF3C, TAFIII90, TFIIB90, mTFIIB90, 2510002F24Rik
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-BRF1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


BRF1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

BRF1 antibody - C-terminal region - Protein Information**BRF1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BRF1 antibody - C-terminal region - Images



WB Suggested Anti-BRF1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Human Thymus

BRF1 antibody - C-terminal region - References

Raineri, I., et al., (2004) (er) Nucleic Acids Res. 32 (4), 1279-1288
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.