

Foxc2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11684**Specification**

Foxc2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q61850
Other Accession	NM_013519 , NP_038547
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Mouse, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

Foxc2 antibody - N-terminal region - Additional Information**Gene ID** 14234**Alias Symbol** Fkh14, Hfhbf3, MFH-1, Mfh1**Other Names**

Forkhead box protein C2, Brain factor 3, BF-3, Forkhead-related protein FKHL14, Mesenchyme fork head protein 1, MFH-1 protein, Transcription factor FKH-14, Foxc2, Fkh14, Fkhl14, Mfh1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Foxc2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Foxc2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Foxc2 antibody - N-terminal region - Protein Information**Name** Foxc2**Synonyms** Fkh14, Fkhl14, Mfh1**Function**

Transcriptional activator.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q99958}.

Tissue Location


Expressed in the brain, muscle, heart, fat, kidney, lung, liver, uterus, ovary and testis.

Foxc2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Foxc2 antibody - N-terminal region - Images



WB Suggested Anti-Foxc2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: Mouse Uterus