

CCBL2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11713**Specification**

CCBL2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A8MY75
Other Accession	NM_001008662 , NP_001008662
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Rabbit, Pig, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

CCBL2 antibody - C-terminal region - Additional Information

Alias Symbol	DKFZp547N1117, DKFZp667D0223, KAT3, MGC9398, RBM1, RBMXL1, RP4-531M19.2, KATIII, RP11-82K18.3
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CCBL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CCBL2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CCBL2 antibody - C-terminal region - Protein Information**CCBL2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCBL2 antibody - C-terminal region - Images



WB Suggested Anti-CCBL2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Human brain

CCBL2 antibody - C-terminal region - References

Yu,P., Gene 365, 111-118 (2006)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.