

U2AF2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11721**Specification**

U2AF2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q80XR5
Other Accession	NM_001012478 , NP_001012496
Reactivity	Human, Mouse, Rat, Zebrafish, Horse, Yeast, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Zebrafish, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 53kDa KDa

U2AF2 antibody - C-terminal region - Additional Information

Alias Symbol **U2AF65**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-U2AF2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


U2AF2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

U2AF2 antibody - C-terminal region - Protein Information**U2AF2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

U2AF2 antibody - C-terminal region - Images



WB Suggested Anti-U2AF2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

There is BioGPS gene expression data showing that U2AF2 is expressed in HepG2