

CARS antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11722**Specification**

CARS antibody - N-terminal region - Product Information

Application	WB
Primary Accession	A8MVQ3
Other Accession	NM_001014438 , NP_001014438
Reactivity	Human, Yeast
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84kDa KDa

CARS antibody - N-terminal region - Additional Information

Alias Symbol **CARS1, CYSRS, MGC:11246**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CARS antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

CARS antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CARS antibody - N-terminal region - Protein Information**CARS antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CARS antibody - N-terminal region - Images



90 kDa
65 kDa
40 kDa
31 kDa
22 kDa

WB Suggested Anti-CARS Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: THP-1 cell lysate

CARS antibody - N-terminal region - References

Liu, C., (2007) J. Mol. Biol. 367 (4), 1063-1078 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.