

Cstf2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11735**Specification**

Cstf2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8BIO5
Other Accession	NM_133196 , NP_573459
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Pig, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 61kDa KDa

Cstf2 antibody - N-terminal region - Additional Information**Gene ID** 108062**Alias Symbol** 64kDa, C630034J23Rik, Cstf64**Other Names**

Cleavage stimulation factor subunit 2, CF-1 64 kDa subunit, Cleavage stimulation factor 64 kDa subunit, CSTF 64 kDa subunit, CstF-64, Cstf2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Cstf2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Cstf2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Cstf2 antibody - N-terminal region - Protein Information**Name** Cstf2**Function**

One of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs. This subunit is directly involved in the binding to pre-mRNAs (By similarity).

Cellular Location

Nucleus. Note=Localized with DDX1 in cleavage bodies.

Tissue Location


Expressed in most somatic cell types (at protein level). Highly expressed in testis, except in meiotic spermatocytes

Cstf2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cstf2 antibody - N-terminal region - Images



WB Suggested Anti-Cstf2 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Pancreas