

EIF4G3 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11753**Specification**

EIF4G3 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O43432-2
Other Accession	NM_001198801 , AAH30578
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine
Predicted	Mouse, Rat, Rabbit, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa

EIF4G3 antibody - N-terminal region - Additional Information

Alias Symbol **eIF4G 3, eIF4GII, eIF-4G 3**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-EIF4G3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


EIF4G3 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

EIF4G3 antibody - N-terminal region - Protein Information**EIF4G3 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EIF4G3 antibody - N-terminal region - Images



70 kDa__
60 kDa__
48 kDa__
36 kDa__
21 kDa__

WB Suggested Anti-EIF4G3 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate

EIF4G3 antibody - N-terminal region - References

Cappelli K, (2006) Res Vet Sci. 2007 Jun;82(3):335-43. Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.