

PCBP4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11824**Specification**

PCBP4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9GZT1
Other Accession	NM_020418 , NP_065151
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Rabbit, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 47kDa KDa

PCBP4 antibody - C-terminal region - Additional Information

Alias Symbol **CBP, LIP4, MCG10**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PCBP4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


PCBP4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PCBP4 antibody - C-terminal region - Protein Information**PCBP4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PCBP4 antibody - C-terminal region - Images



WB Suggested Anti-PCBP4 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate

PCBP4 antibody - C-terminal region - References

Chkheidze, A.N. (2003) Mol. Cell. Biol. 23 (23), 8405-8415
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.