

**Lsm10 Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI11839****Specification**

---

**Lsm10 Antibody - N-terminal region - Product Information**

|                   |                                                             |
|-------------------|-------------------------------------------------------------|
| Application       | WB                                                          |
| Primary Accession | <a href="#">B2RYU1</a>                                      |
| Other Accession   | <a href="#">NM_001108976</a> , <a href="#">NP_001102446</a> |
| Reactivity        | Human, Mouse, Rat, Rabbit, Zebrafish, Horse, Bovine, Dog    |
| Predicted         | Human, Mouse, Rat, Pig, Horse, Bovine, Dog                  |
| Host              | Rabbit                                                      |
| Clonality         | Polyclonal                                                  |
| Calculated MW     | 13kDa KDa                                                   |

**Lsm10 Antibody - N-terminal region - Additional Information****Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Lsm10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Lsm10 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


**Lsm10 Antibody - N-terminal region - Protein Information****Lsm10 Antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Lsm10 Antibody - N-terminal region - Images**

---



Host: Rabbit  
Target Name: Lsm10  
Sample Tissue: Rat Testis lysates  
Antibody Dilution: 1.0µg/ml