

EIF4E3 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11852**Specification**

EIF4E3 antibody - middle region - Product Information

Application	WB
Primary Accession	Q8N5X7
Other Accession	NM_001134649 , NP_001128123
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Rat, Rabbit, Zebrafish, Chicken, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 24kDa KDa

EIF4E3 antibody - middle region - Additional Information**Gene ID** 317649**Alias Symbol** MGC39820, MGC86971**Other Names**

Eukaryotic translation initiation factor 4E type 3, eIF-4E type 3, eIF-4E3, eIF4E type 3, eIF4E-3, EIF4E3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-EIF4E3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

EIF4E3 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

EIF4E3 antibody - middle region - Protein Information**Name** EIF4E3**Function**


Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation of protein synthesis. May act as an inhibitor of EIF4E1 activity (By similarity).

EIF4E3 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EIF4E3 antibody - middle region - Images



WB Suggested Anti-EIF4E3 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Human Liver