

NKD2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11870**Specification**

NKD2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O96EK8
Other Accession	NM_033120 , AAH12176
Reactivity	Human, Mouse, Rat, Horse
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

NKD2 antibody - N-terminal region - Additional Information

Alias Symbol	Naked2
Format	

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NKD2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


NKD2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NKD2 antibody - N-terminal region - Protein Information**NKD2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NKD2 antibody - N-terminal region - Images



WB Suggested Anti-NKD2 Antibody Titration: 0.2-1 $\mu\text{g/ml}$
Positive Control: Jurkat cell lysate

NKD2 antibody - N-terminal region - References

Strausberg, R. Direct Submission Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.