

Wdr12 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11875**Specification**

Wdr12 antibody - middle region - Product Information

Application	WB
Primary Accession	O9JJA4
Other Accession	NM_021312 , NP_067287
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Yeast, Bovine, Dog
Predicted Host	Mouse, Rat, Zebrafish, Chicken, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 47kDa KDa

Wdr12 antibody - middle region - Additional Information**Gene ID** 57750**Alias Symbol** 4933402C23Rik, Ytm1, Ytm1p**Other Names**

Ribosome biogenesis protein WDR12 {ECO:0000255|HAMAP-Rule:MF_03029}, WD repeat-containing protein 12 {ECO:0000255|HAMAP-Rule:MF_03029}, Wdr12

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Wdr12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Wdr12 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Wdr12 antibody - middle region - Protein Information**Name** Wdr12**Function**

Component of the PeBoW complex, which is required for maturation of 28S and 5.8S ribosomal RNAs and formation of the 60S ribosome.

Cellular Location


Nucleus, nucleolus {ECO:0000255|HAMAP-Rule:MF_03029}. Nucleus, nucleoplasm {ECO:0000255|HAMAP-Rule:MF_03029}

Wdr12 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Wdr12 antibody - middle region - Images



WB Suggested Anti-Wdr12 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Spleen