

SLC12A1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11886**Specification**

SLC12A1 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q8IUN5
Other Accession	NM_000338 , AAH40138
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47kDa KDa

SLC12A1 antibody - N-terminal region - Additional Information

Alias Symbol **BSC1, MGC48843, NKCC2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SLC12A1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


SLC12A1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLC12A1 antibody - N-terminal region - Protein Information**SLC12A1 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SLC12A1 antibody - N-terminal region - Images



Rabbit Anti-SLC12A1 Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-SLC12A1 Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate



Host: Rabbit
Target Name: SLC12A1
Sample Tissue: Jurkat
Lane A: Primary Antibody
Lane B: Primary Antibody + Blocking Peptide
Primary Antibody
Concentration: 1 µg/ml
Peptide Concentration: 5 µg/ml
Lysate Quantity: 25 µg/lane
Gel Concentration: 0.12

