

GGTLA4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11919**Specification**

GGTLA4 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q9BX51
Other Accession	NM_080920 , NP_563577
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted Host	Human
Clonality	Rabbit
Calculated MW	Polyclonal 25kDa KDa

GGTLA4 antibody - C-terminal region - Additional Information**Gene ID** 92086**Alias Symbol** RP5-831C21.2, MGC50550, dJ831C21.2, GGTL6, GGTLA3, GGTLA4, dJ831C21.1**Other Names**

Gamma-glutamyltransferase light chain 1, Gamma-glutamyltransferase-like activity 4, Gamma-glutamyltransferase-like protein 6, GTTLC1, GGTLA4

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-GGTLA4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


GGTLA4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GGTLA4 antibody - C-terminal region - Protein Information**Name** GTTLC1**Synonyms** GGTLA4**GGTLA4 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GGTLA4 antibody - C-terminal region - Images



Rabbit Anti-GGTLA4 Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-GGTLA4 Antibody Titration: 5.0 µg/ml
Positive Control: Human kidney