

Stc1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11977**Specification**

Stc1 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P97574
Other Accession	NM_031123 , NP_112385
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27kDa KDa

Stc1 Antibody - C-terminal region - Additional Information**Gene ID** 81801**Other Names**

Stanniocalcin-1, STC-1, Stc1, Stc

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Stc1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Stc1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Stc1 Antibody - C-terminal region - Protein Information**Name** Stc1**Synonyms** Stc**Function**

Stimulates renal phosphate reabsorption, and could therefore prevent hypercalcemia.

Cellular Location


Secreted.

Stc1 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Stc1 Antibody - C-terminal region - Images



WB Suggested Anti-Stc1 Antibody Titration: 1.0 µg/ml
Positive Control: Rat Pancreas