

SLC17A3 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11996**Specification**

SLC17A3 antibody - middle region - Product Information

Application	WB
Primary Accession	O9H533
Other Accession	NM_006632 , NP_006623
Reactivity	Human, Horse, Bovine
Predicted	Human, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46kDa KDa

SLC17A3 antibody - middle region - Additional Information

Alias Symbol	NPT4
--------------	------

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SLC17A3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


SLC17A3 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

SLC17A3 antibody - middle region - Protein Information**SLC17A3 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC17A3 antibody - middle region - Images



WB Suggested Anti-SLC17A3 Antibody Titration: 0.2-1 μ g/ml
Positive Control: HepG2 cell lysate

SLC17A3 antibody - middle region - References

Melis,D., (2004) J. Inherit. Metab. Dis. 27 (6), 725-733
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.