

PSG1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12002**Specification**

PSG1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6ICR4
Other Accession	NM_006905 , NP_008836
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47kDa KDa

PSG1 antibody - C-terminal region - Additional Information

Alias Symbol	B1G1, CD66f, DHFRP2, FLJ90598, FLJ90654, PBG1, PSBG1, PSG IIA-d, PSGGA, PSGIIA-a, PSGIIA-b, PSGIIA-c, SP1, PSGIIA
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-PSG1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


PSG1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PSG1 antibody - C-terminal region - Protein Information**PSG1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PSG1 antibody - C-terminal region - Images



WB Suggested Anti-PSG1 Antibody Titration: 2.5µg/ml

Positive Control: NCI-H226 cell lysate

.PSG1 is strongly supported by BioGPS gene expression data to be expressed in Human NCI-H226 cells

PSG1 antibody - C-terminal region - References

Otsuki,T., (2005) DNA Res. 12 (2), 117-126
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.