

SERPINB13 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12007**Specification**

SERPINB13 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9UIV8
Other Accession	NM_012397 , NP_036529
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted Host	Human, Horse, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 44kDa KDa

SERPINB13 antibody - N-terminal region - Additional Information**Gene ID** 5275

Alias Symbol	HUR7, MGC126870, PI13, headpin, HSHUR7SEQ
--------------	---

Other Names

Serpin B13, HaCaT UV-repressible serpin, Hurpin, Headpin, Peptidase inhibitor 13, PI-13, Proteinase inhibitor 13, SERPINB13, PI13

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SERPINB13 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SERPINB13 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SERPINB13 antibody - N-terminal region - Protein Information

Name SERPINB13

Synonyms PI13

Function

May play a role in the proliferation or differentiation of keratinocytes.

Cellular Location

Cytoplasm.


Tissue Location

Skin specific.

SERPINB13 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SERPINB13 antibody - N-terminal region - Images

WB Suggested Anti-SERPINB13 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: COLO205 cell lysate

There is BioGPS gene expression data showing that SERPINB13 is expressed in COLO205

SERPINB13 antibody - N-terminal region - References

Bylaite, M., (2006) Exp. Dermatol. 15 (2), 110-118 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.