

ZNF17 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12039**Specification**

ZNF17 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8N7M1
Other Accession	NM_006959 , NP_008890
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

ZNF17 antibody - N-terminal region - Additional Information

Alias Symbol **HPF3, KOX10**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF17 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ZNF17 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF17 antibody - N-terminal region - Protein Information**ZNF17 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF17 antibody - N-terminal region - Images



WB Suggested Anti-ZNF17 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:12500

Positive Control: Human Small Intestine

ZNF17 antibody - N-terminal region - References

Huebner, K., et al., (1991) Am. J. Hum. Genet. 48(4), 726-740 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.