

KIAA1618 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12041**Specification**

KIAA1618 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9HCF4-2
Other Accession	NM_020954 , NP_066005
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	118kDa KDa

KIAA1618 antibody - N-terminal region - Additional Information

Alias Symbol	ALO17, MYMY2, MYSTR, NET57, C17orf27, KIAA1618
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-KIAA1618 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


KIAA1618 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

KIAA1618 antibody - N-terminal region - Protein Information**KIAA1618 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KIAA1618 antibody - N-terminal region - Images



WB Suggested Anti-KIAA1618 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Hela cell lysate

KIAA1618 antibody - N-terminal region - References

Cools,J.,(2002)GenesChromosomesCancer34(4),354-362ReconstitutionandStorage:Forshorttermuse ,storeat2-8Cupto1week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.