

WT1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12067**Specification**

WT1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	E9PMK7
Other Accession	NM_000378 , NP_000369
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

WT1 antibody - N-terminal region - Additional Information

Alias Symbol	AWT1, GUD, WAGR, WIT-2, WT33, NPHS4, EWS-WT1
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-WT1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

WT1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

WT1 antibody - N-terminal region - Protein Information**WT1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WT1 antibody - N-terminal region - Images



WB Suggested Anti-WT1 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: Hela cell lysate